

Corn - Post Herbicide Performance

Herbicides	B. Nightshade	Velvetleaf	Waterhemp	R. Pigweed	Cocklebur	Lambsquarter	W. Buckwheat	Russian Thistle	Jimsonweed	Kochia	Giant Ragweed	Common Ragweed	Penn. Smartweed	Morningglory	Palmer Amaranth	Sunflower
Aim EW 0.5 oz	F	F	G	F	G	F	F	G	F/G	F/G	F	F	G	G/E	F	F
Glyphosate 32 oz	G/E	G	F	F	F	G	F	G	F	F	G/E	G/E	G	F	G/E	F
Aim EW 0.5 oz + Glyphosate 32 oz	F	F	F	F	F	F	F	G	G	F	F	F	F	F	F	F
Aim EW 0.5 oz + Atrazine 1/2 lb	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Atrazine 1/2 lb + Clarity® 4 oz	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Clarity® 8 oz	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Hornet®WDG 2.5-3 oz	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Distinct® 4.0 oz	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Callisto® 1.5 oz + Atrazine 1/2 lb	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Equip® 1.25 oz	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Liberty® 32 oz	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Starane® 0.5 pt + 1/2 lb Atrazine	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Steadfast® 3/4 oz + 1/2 lb Atrazine	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Aim EW 0.5 oz + Steadfast® 3/4 oz + 4 oz Clarity®	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F

F = Fair Control **G** = Good Control **E** = Excellent Control

Adjuvant Guidelines

- COC at 1 gal/100 gals or NIS at 1 qt/100 gals of spray solution (or equivalent product).
- Use of MSO is allowed where required by tankmix partner.
- Follow tankmix partner(s) label for all adjuvants recommended. (AMS/UAN).

Application Guidelines

- 15-20 GPA with flat fan type nozzles recommended.
- Coverage is essential for good control.
- Broadcast spray up to 8 leaf collar corn (V8).
- Directed spray with drop nozzles from 8-14 leaf corn.

Nozzle Selection Guidelines

- For best control use tips rated as excellent for contact herbicides.
- Tips rated as good for contact herbicides will require higher gals/psi.
- Do not use tips rated as poor for contact herbicides.
- Do not use flood type nozzles, course spray, for foliar treatments.

Avoiding Crop Response

- Apply when crop is actively growing, allowing rapid metabolism of herbicide.
- Do not make applications during cool, damp (adverse) weather conditions.
- Cool conditions (averaging less than 65° F) along with high humidity (70%+) and/or wet leaves (dew, rain, irrigation) can increase symptoms of crop response.
- Applications should not be made within 6-8 hours of rainfall/irrigation to avoid movement into the whorl and concentrating product.
- Aim herbicide is a contact product only. No systemic movement will occur in the plant.

Aim Post Corn Value

- Premium Velvetleaf Control.
- Improves control of:
 - Waterhemp
 - Lambsquarter
 - Nightshade
 - Morningglories
 - Kochia
 - Russian thistle
- Fast removal of weed competition.
- Wide application window.

